


Distribution of the endangered red-tailed black-cockatoo (south-eastern) *(Calyptorhynchus banksii graptogyne)*

This map shows the general distribution of *Calyptorhynchus banksii graptogyne*. The known species and breeding distribution comprises confirmed observation points buffered by 1km that occurred within mapped vegetation (NVIS 4.1) of important breeding and foraging tree species. The likely and may occur distributions were defined using statistical modelling (Maxent) and the intersection with vegetation mapping of important feed and nesting tree species. Observation data provided by the Victorian Biodiversity Atlas (DEPI), Biological Databases of South Australia (DEWNR), Birdlife Australia, Australian National Wildlife Collection (CSIRO), Australian Bird and Bat Banding Scheme (DotE) and state museums. Data was also provided by Richard Hill of the Victorian Dept of Environment and Primary Industries. Distribution has been generalised to 10km. Distribution mapped by ERIN Nov 2006, updated Feb 2016.

- Species known to occur
- Species likely to occur
- Species may occur



Produced by:
Environmental Resources Information Network
Australian Government
Department of the Environment and Energy
July 2016
© Commonwealth of Australia, 2016

Datum: Geocentric Datum of Australia 1994

Caveat: The information in this map has been compiled from a range of sources. While every effort has been made to ensure accuracy and completeness, no guarantee is given, nor responsibility taken by the Commonwealth for errors or omissions, and the Commonwealth does not accept responsibility in respect of any information or advice given in relation to, or as a consequence of, anything containing herein.